

Multifold Label
LEAFLET

**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

**BIVATOP 200
Long Acting Injectable**

Composition

Each mL contains 200 mg oxytetracycline (as dihydrate).

Indications

For the treatment of conditions caused by, or associated with oxytetracycline-sensitive organisms and infections in cattle, sheep and pigs.

Directions for Use

Restraints:

DO NOT administer more than two injections to any pig.

The use of tetracycline during the period of tooth development, including late pregnancy, may lead to tooth discolouration. Occasional sensitivity reactions (anaphylaxis) to oxytetracycline have been known to occur, administer adrenaline immediately.

Dosage and Administration:

Discard 90 days after initial use when stored below 25°C (Air Conditioning).

Administer 1 mL per 10 kg bodyweight by subcutaneous injection in cattle or by deep intramuscular injection in cattle, sheep or pigs to provide a dosage of 20 mg oxytetracycline per kg bodyweight.

If necessary a second dose may be administered 72 hours after the initial dose. The maximum recommended dose volume is 10 mL for intramuscular and subcutaneous injection.

If given intravenously a second dose may be administered 12-24 hours after the initial dose.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period. Injections should be given in the neck region. Injection volumes should not exceed 10 mL per site. Treated animals should be clearly identified in such a way that they will maintain their identity during the withholding period.

WITHHOLDING PERIOD:

MEAT (CATTLE): DO NOT USE less than 21 days before slaughter for human consumption.

MEAT (SHEEP): DO NOT USE less than 42 days before slaughter for human consumption.

MEAT (PIGS): DO NOT USE less than 21 days before slaughter for human consumption.

MILK (CATTLE/SHEEP): Milk collected from cows or ewes within 7 days following treatment MUST NOT BE USED for human consumption or processing, or fed to bobby calves.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (PIGS): DO NOT slaughter for export less than 21 days after final treatment.

EXPORT SLAUGHTER INTERVAL (CATTLE/SHEEP): This product does not have an ESI established. For advice on the ESI, contact the Manufacturer on 1800 038 037 before using this product.

The ESI on this label was current at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1800 038 037 or the APVMA website.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre, phone Australia 13 11 26.

Disposal

Dispose of empty container by wrapping with paper and putting in garbage.

Storage

Store below 25°C (Air Conditioning).

Presentation

100 or 250 mL amber glass injection vials.

Distributed by
Boehringer Ingelheim Pty Limited
Animal Health Division
78 Waterloo Road
NORTH RYDE NSW 2113

APVMA 51637/56736

Bivatop® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH – used under license.