

**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

Enterisol® Ileitis

attenuated live vaccine



Composition

One 2 mL dose of reconstituted vaccine contains (titres in TCID₅₀):

Lawsonia intracellularis

At least 1 x 10^{4.9}

* Tissue Culture Infective Dose

Indications

For active immunisation of pigs from three weeks of age and older as an aid in the prevention and control of porcine proliferative enteropathy (Ileitis) caused by *Lawsonia intracellularis* infection.

DIRECTIONS FOR USE

DO NOT vaccinate breeding boars.

DO NOT mix with other vaccine/immunological product, antibiotics, or antimicrobial products.

Antibiotics should not be administered from 3 days prior to vaccination until 3 days post vaccination.

Reconstitute the vaccine by adding the full content of the accompanying diluent. Shake well and use immediately.

Vaccination by oral drenching:
Administer a single 2 mL dose orally, irrespective of bodyweight.

Vaccination via the drinking water:
The systems have to be cleaned and rinsed with untreated water to avoid any residues of chemotherapeutics, detergents or disinfectants.

The final solution containing the vaccine should be consumed within four hours after preparation. Calculate the number of vials required to vaccinate all the pigs according to the table below.

No. of pigs	Vaccine vial	Diluent vial
10	10 dose (20 mL)	20 mL
50	50 dose (100 mL)	100 mL
100	100 dose (100 mL)	200 mL

Dilute the vaccine in drinking water on the basis of pre-measured water intake during a four hour time period of the previous day at the time of planned vaccination.

It is recommended to add skimmed milk powder as a stabilizer into the drinking water prior to adding the vaccine. The final concentration of the skimmed milk should be 2.5 g/litre.

A clinical disease history or serology in grower pigs is recommended prior to vaccination.

WITHHOLDING PERIOD: NIL

Disposal

Dispose of empty container by wrapping in paper and putting in garbage.

Storage

Store between 2°C and 8°C. Do not freeze. Use entire contents when first opened.

Packaging

Vaccine: Amber glass vial of 20 mL (10 doses), 100 mL (50/100 doses), and 12 x 100 mL (12 x 100 doses).

Diluent: Clear glass vial containing 20 mL (10 doses), 100 mL (50 doses), 200 mL (100 doses), and 12 x 200 mL (12 x 100 doses).
High density polyethylene vial containing 20 mL (10 doses), 100 mL (50 doses), 200 mL (100 doses), and 12 x 200 mL (12 x 100 doses).

Australia

**Boehringer Ingelheim Pty Limited
Animal Health Division
78 Waterloo Road
North Ryde, N.S.W. 2113
APVMA 58972/0306**

New Zealand

**Boehringer Ingelheim (NZ) Limited
Animal Health Division
Level 1, Unit 9
42 Ormiston Road
East Tamaki, Auckland**

Restricted Veterinary Medicine. A9569

See www.foodsafety.govt.nz for registration conditions.

Enterisol® is a registered trademark of
Boehringer Ingelheim Vetmedica GmbH, used
under license.

Boehringer Ingelheim

Revision 6
Operator lsh
Date 04Dec13
Art # 130444-03
Item # 430438
SAP #
Product Enterisol Ileitis
Type/Size Insert
Template # 671
Dimensions 5-5/8" x 8-3/4"
Fold 1-7/8" x 1-7/8"
PDF 04Dec13 lsh
Rev Level 03

BLACK

**Tradename - 18 pt.
Text - 8.7 pt.**

Art # - 7 pt.