

**READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY****Fel-O-Vax® 5**  **Boehringer
Ingelheim**
**Vaccine
(Inactivated)****Active constituents**

Each 1 mL contains inactivated feline rhinotracheitis virus (FVR 605, RP ≥ 1.0), feline calicivirus (FCV 255, RP ≥ 1.0), panleucopenia virus (FPV CU-4, RP ≥ 1.0), *Chlamydomphila felis* (CELLO, RP ≥ 1.0) and feline leukaemia virus (1161E, RP ≥ 1.0), amphotericin, polymyxin B, neomycin and thiomersal (<0.05% w/v) added as preservatives.

An inactivated vaccine for the active immunisation of healthy cats 8 weeks of age or older to aid in the reduction of clinical symptoms caused by infection with feline rhinotracheitis virus, feline calicivirus, panleucopenia virus, *Chlamydomphila felis* and feline leukaemia virus.

This vaccine does not prevent infection or shedding.

To be used by, or under the direction of a registered veterinary surgeon.

Net contents 25 x 1 mL vials

Directions for Use**Precautions**

Vaccinate only healthy cats or kittens with normal temperatures.

Cats under treatment with immunosuppressive drugs should be vaccinated with caution.

Studies to support the use of this product in pregnant queens have not been conducted.

Side effects

Some animals may show transient post-vaccination reactions including pain, malaise, lethargy, inappetence, transient fever, depression and/or stiffness. These reactions usually disappear within 24-48 hours.

Allergic reactions such as anaphylaxis occur rarely and may require parenteral treatment with an antihistamine, a corticosteroid or adrenalin as appropriate.

Local site reactions after vaccination are rare but transient swelling, pain and subcutaneous thickening have been reported.

Based on overseas challenge data, it has been demonstrated that clinical signs of feline calicivirus infection (pyrexia, ulcers) are likely to occur in vaccinated animals following exposure to this virus. However, the severity of the fever and ulcers is likely to be reduced compared to unvaccinated animals.

Dosage and administration

Shake the bottle well to obtain a uniform suspension of the vaccine before administration.

The entire contents of the bottle should be used immediately after opening.

A new sterile syringe and needle should always be used to administer this vaccine.

Dose

Withdraw the contents of a single dose vial (1 mL) aseptically into a sterile syringe and inject subcutaneously.

Recommended vaccination plan

Fel-O-Vax 5 vaccine can be used on its own or as part of a standard vaccination program.

The following vaccination program is recommended for cats and kittens.

Cats aged 8 weeks and older	For an initial vaccination course, two doses are required, injected at an interval of 3-4 weeks
Booster Vaccination	Annually boosting with a single dose of vaccine is recommended

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Additional user safety information

Self-Injection: Take care to avoid self-injection. In the event of self-administration, seek prompt medical attention and take this package leaflet/carton with you. If possible the application of gentle squeezing pressure with absorbent material (e.g. facial tissues) at the injection site will swab up unabsorbed vaccine. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleansed and a topical antiseptic applied. Accidental self-administration may result in local bruising, severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If pain persists for more than 12 hours after medical examination, seek medical advice again.

Disposal

Discarded syringes and needles should immediately be placed in a designated and appropriately labelled "sharps" container. Discarded vials should immediately be placed in a designated and appropriately labelled "biologicals" container.

Storage

Store between 2-8°C. Do not freeze. Protect from light.

**Boehringer Ingelheim Pty Limited
Animal Health Division**

78 Waterloo Road
NORTH RYDE NSW 2113
Phone 1800 038 037
www.boehringer-ingelheim.com.au
APVMA Approval No. 51366/57811

Manufactured by
Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506 USA
Made in USA

Packaging

Vaccines are supplied as 25 or 10 vials of inactivated vaccine.

Not all pack sizes may be marketed.

It is illegal to sell individual vials separately.

Fel-O-Vax® is a registered trademark of Boehringer Ingelheim Vetmedica, Inc. - used under license.



Please recycle leaflet, label and tray only.

Boehringer Ingelheim

Revision 5
Operator lsh
Date 16Sep16
Art # 154801-01
SAP # 108387-002
Product Fel-O-Vax 5
Type/Size Insert
Template # 544
Dimensions 3" x 11"
Fold 3" x 1"
PDF 16Sep16 lsh

BLACK

This file is 100% when printed on A4 (or A3, as applicable)

DIMENSIONS: 76.2 mm x 279.4 mm

FONT SIZES:

Tradename: 16 pt

Text: 9 pt

Art Number: 7 pt