

**READ SAFETY DIRECTIONS  
FOR ANIMAL TREATMENT ONLY**
**Fel-O-Vax® 5 Vaccine (Inactivated)**

An inactivated vaccine for the active immunisation of healthy cats 8 weeks of age or older to aid in the reduction of clinical symptoms caused by infection with feline rhinotracheitis virus, feline calicivirus, panleucopenia virus, *Chlamydomphila felis* and feline leukaemia virus.

This vaccine does not prevent infection or shedding.

**Composition:**

Each 1 mL contains inactivated feline rhinotracheitis virus (FVR 605, RP ≥ 1.0), feline calicivirus (FCV 255, RP ≥ 1.0), panleucopenia virus (FPV CU-4, RP ≥ 1.0), *Chlamydomphila felis* (CELLO, RP ≥ 1.0) and feline leukaemia virus (1161E, RP ≥ 1.0), amphotericin, polymyxin B, neomycin and thiomersal (<0.05% w/v) added as preservatives.

**To be used by, or under the direction of a registered veterinary surgeon.**

**DIRECTIONS FOR USE**
**Precautions:**

**Vaccinate only healthy cats or kittens with normal temperatures.**

**Cats under treatment with immunosuppressive drugs should be vaccinated with caution.**

**Studies to support the use of this product in pregnant queens have not been conducted.**

**Side effects:**

Some animals may show transient post-vaccination reactions including pain, malaise, lethargy, inappetence, transient fever, depression and/or stiffness. These reactions usually disappear within 24-48 hours.

Allergic reactions such as anaphylaxis occur rarely and may require parenteral treatment with an antihistamine, a corticosteroid or adrenalin as appropriate.

Local site reactions after vaccination are rare but transient swelling, pain and subcutaneous thickening have been reported.

Based on overseas challenge data, it has been demonstrated that clinical signs of feline calicivirus infection (pyrexia, ulcers) are likely to occur in vaccinated animals following exposure to this virus. However, the severity of the fever and ulcers is likely to be reduced compared to unvaccinated animals. This vaccine does not prevent infection or shedding.

**Dosage and Administration:**

**Shake the bottle well to obtain a uniform suspension of the vaccine before administration.**

**The entire contents of the bottle should be used immediately after opening.**

**A new sterile syringe and needle should always be used to administer this vaccine.**

**Dose:**

Withdraw the contents of a single dose vial (1 mL) aseptically into a sterile syringe and inject subcutaneously.

**Recommended Vaccination Plan:**

Fel-O-Vax 5 vaccine can be used on its own or as part of a standard vaccination program.

The following vaccination program is recommended for cats and kittens.

Cats aged 8 weeks and older	For an initial vaccination course, two doses are required, injected at an interval of 3-4 weeks
Booster Vaccination	Annually boosting with a single dose of vaccine is recommended

**User Safety Information:**

**Self-Injection:** Take care to avoid self-injection. In the event of self-administration, seek prompt medical attention and take this package leaflet/carton with you. If possible the application of gentle squeezing pressure with absorbent material (e.g. facial tissues) at the injection site will swab up unabsorbed vaccine. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleansed and a topical antiseptic applied. Accidental self-administration may result in local bruising, severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If pain persists for more than 12 hours after medical examination, seek medical advice again.

**First Aid:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia *131126*.

**Disposal:**

Discarded syringes and needles should immediately be placed in a designated and appropriately labelled "sharps" container. Discarded vials should immediately be placed in a designated and appropriately labelled "biologicals" container.

**Storage:**

Store between 2-8°C. Do not freeze. Protect from light.

**Packaging:**

Vaccines are supplied as 25 or 10 vials of inactivated vaccine.

Not all pack sizes may be marketed.

Fel-O-Vax® 5 is a registered trademark of Boehringer Ingelheim Vetmedica, Inc. - used under license.

APVMA Approval No. 51366/57811

**Boehringer Ingelheim Pty Limited**

**Animal Health Division  
78 Waterloo Road  
NORTH RYDE NSW 2113**



Please recycle leaflet, label and tray only.

**Boehringer Ingelheim**

Revision ..... 11  
Operator ..... Ish  
Date ..... 13Nov13  
Art # ..... 154801-00  
Item # ..... 454802  
SAP # .....  
Product ..... Fel-O-Vax 5  
Type/Size ..... Insert  
Template # ... 537  
Dimensions... 3" x 8-1/2"  
Fold ..... 3" x 1-1/16"  
PDF ..... 13Nov13 Ish  
Rev Level ..... 00

**BLACK**

This file is 100% when printed on A4 (or A3, as applicable)

**DIMENSIONS:** 76.2 mm x 215.9 mm

**FONT SIZES:**

Tradename: 12 pt

Art Number: 7 pt

Text: 8.5 pt