

1309147-02

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Ingelvac CircoFLEX®

Suspension for Injection in Pigs

Each 1 mL dose contains:

Porcine circovirus type 2 ORF2 protein, ≥ 1.0 RP

Carbomer 1 mg

Indications

For active immunisation of pigs from two weeks of age and older as an aid in the prevention and control of diseases associated with porcine circovirus type 2. This vaccine has been demonstrated safe in pregnant and breeding sows.

Net contents

10 mL, 50 mL, 100 mL, or 250 mL

DIRECTIONS FOR USE

SHAKE WELL BEFORE USE and use immediately.

Use the entire contents after broaching the container. Administer a single 1 mL dose intramuscularly, irrespective of bodyweight.

Precautions

Interaction: No information is available on the safety and efficacy of the concurrent use of this vaccine with any other products except with Boehringer Ingelheim's Ingelvac MycoFLEX®.

Side effects

A mild and transient hyperthermia very commonly occurs on the day of vaccination.

Dosage and administration


Administer a single 1 mL dose per pig intramuscularly, irrespective of body weight.

When mixed with Ingelvac MycoFLEX®:

- Vaccinate only pigs from 3 weeks of age.
- Use the same volumes of Ingelvac CircoFLEX® and Ingelvac MycoFLEX®.

To ensure correct mixing follow the steps as described below:

- Use a pre-sterilised transfer needle.
Pre-sterilised transfer needles are commonly available via medical equipment suppliers.
- Connect one end of the transfer needle to the vaccine bottle of Ingelvac MycoFLEX®.
- Connect the opposite end of the transfer needle to the vaccine bottle of Ingelvac CircoFLEX®. Transfer the Ingelvac CircoFLEX® vaccine into the vaccine bottle of Ingelvac MycoFLEX®. If needed, gently press the vaccine bottle of Ingelvac CircoFLEX® to facilitate the transfer. After the transfer of the full content of Ingelvac CircoFLEX®, disconnect the transfer needle. Discard transfer needle and the empty vaccine bottle of Ingelvac CircoFLEX®.
- To ensure appropriate mixing of the vaccines, gently shake the vaccine bottle until the mixture is of uniform orange to reddish colour. During vaccination the uniformity of the coloured mixture should be monitored and maintained by continuous agitation.
- Administer one single injection dose (2 mL) of the mixture intramuscularly per pig, irrespective of body weight. For administration, vaccine devices should be used in accordance with the device instructions provided by the manufacturer.



- Use the entire vaccine mixture immediately after mixing.



The package leaflet of Ingelvac MycoFLEX® should also be consulted before the administration of the mixed product.

General directions

Caution: avoid carcass damage

1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use needles of appropriate gauge and length.
5. As far as possible, avoid injection of animals during wet weather or under dusty conditions.
6. This product should be injected only into muscle tissue.

WITHHOLDING PERIOD

Zero (0) days

TRADE ADVICE - Export Slaughter Interval (ESI) :

Zero (0) days

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Additional user safety information

Take care to avoid self-injection. Accidental self-administration may result in local bruising, pain and swelling. In the event of self-administration, seek medical attention and show the package leaflet or label to the Medical Practitioner.

Disposal

Dispose of empty container by wrapping in paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.

Storage

Store between 2°C - 8°C.

Do not freeze.

Protect from light.

Packaging

High-density polyethylene bottles of 10 mL, 50 mL, 100 mL or 250 mL with a chlorobutyl stopper and lacquered aluminium cover.

APVMA: 63488/106519

Australia
Boehringer Ingelheim Pty Limited
Animal Health Division
78 Waterloo Road
North Ryde NSW 2113
Phone: 1800 038 037
www.boehringer-ingelheim.com.au



Boehringer Ingelheim

Revision. 1
Operator. hal
Date. 03Mar17
Art #. 1309147-02
SAP #. 106685-003
Product. Ingelvac CircoFLEX
Type/Size. Insert
Template #. 671
Dimensions. 5-5/8" x 8-3/4"
Folds To. 1-7/8" x 1-1/8"
PDF. 03Mar17 hal

BLACK

Tradename: 22 pt.

Body: 7.4 pt.