

130961-02

**KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE**

Ingelvac CircoFLEX[®]



suspension for injection for pigs

Active Constituent

Each 1 mL dose contains:
Porcine circovirus type 2 ORF2 protein
Min RP* 1.0, Max RP* 3.75
* Relative Potency

For active immunisation of pigs from two weeks of age and older as an aid in the prevention and control of porcine circovirus type 2 associated diseases caused by porcine circovirus type 2.

DIRECTIONS FOR USE

DO NOT mix with other vaccine/immunological products except for Ingelvac MycoFLEX[®].
Shake well and use immediately.
Administer a single 1 mL dose intramuscularly, irrespective of body weight.

To be mixed with Ingelvac MycoFLEX[®]:

For immunisation from three weeks of age.
Use the same package size of each vaccine.
Aseptically transfer the full content of each vaccine bottle into a pre-sterilised plastic or glass container of at least double the capacity of the vaccine bottle by using pre-sterilised transfer needles. Shake well until a uniform colour of the mixture is obtained.

Administer one dose (2 mL) of the mixture intramuscularly to each pig. Use entire content immediately after mixing.

WITHHOLDING PERIOD: NIL

Disposal

Dispose of empty container by wrapping in paper and putting in garbage.

Storage

Store between 2°C and 8°C.
Do not freeze. Protect from light.
Use entire contents when first opened.

Packaging

High-density polyethylene bottles of 10 mL (10 doses), 50 mL (50 doses), 100 mL (100 doses), or 250 mL (250 doses) with a chlorobutyl stopper and lacquered aluminium seal.

Restricted Veterinary Medicine A9935

See www.foodsafety.govt.nz for registration conditions.

Ingelvac CircoFLEX[®] is a registered trademark of Boehringer Ingelheim Vetmedica GmbH. Used under license.

Distributed by

**Boehringer Ingelheim (NZ) Limited
Animal Health Division
Level 1, Unit 9
42 Ormiston Road
East Tamaki, Auckland**

