

**KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

Ingelvac MycoFLEX®

Mycoplasma hyopneumoniae inactivated vaccine for pigs



Active Constituent

Each 1 mL dose contains:

Inactivated *Mycoplasma hyopneumoniae* ≥ 1.0 RP
Carbomer 1mg

Indications

For the vaccination of healthy, susceptible swine at weaning (3 weeks of age) or older as an aid in the prevention of enzootic pneumonia of swine caused by *Mycoplasma hyopneumoniae*.

DIRECTIONS FOR USE

Restraints

DO NOT mix with other vaccine/immunological products, except with Boehringer Ingelheim's Ingelvac CircoFLEX.

DO NOT administer during pregnancy and lactation as no safety or efficacy data has been generated in breeding animals.

Side effects

A mild and transient hyperthermia very commonly occurs on the day of vaccination.

Dosage and Administration

SHAKE WELL BEFORE USE and use immediately.

During use, hold vaccine at or below 25°C.

Use entire contents after broaching the container.

Administer a single 1 mL dose per pig intramuscularly.


When mixed with Ingelvac CircoFLEX®:

- Vaccinate only pigs from 3 weeks of age.
- Use the same volumes of Ingelvac CircoFLEX and Ingelvac MycoFLEX.
- Use a pre-sterilised transfer needle. Pre-sterilised transfer needles are commonly available via medical equipment suppliers.

To ensure correct mixing follow the steps as described below:

- Connect one end of the transfer needle to the vaccine bottle of Ingelvac MycoFLEX.
- Connect the opposite end of the transfer needle to the vaccine bottle of Ingelvac CircoFLEX. Transfer the Ingelvac CircoFLEX vaccine into the vaccine bottle of Ingelvac MycoFLEX. If needed, gently press the vaccine bottle of Ingelvac CircoFLEX to facilitate the transfer. After the transfer of the full content of Ingelvac CircoFLEX, disconnect the transfer needle. Discard transfer needle and the empty vaccine bottle of Ingelvac CircoFLEX.
- To ensure appropriate mixing of the vaccines, gently shake the vaccine bottle of Ingelvac MycoFLEX until the mixture is of uniform orange to reddish colour. During vaccination the uniformity of the coloured mixture should be monitored and maintained by continuous agitation.

- Administer one single injection dose (2 mL) of the mixture intramuscularly per pig, irrespective of body weight. For administration, vaccine devices should be used in accordance with the device instructions provided by the manufacturer.



Use the entire vaccine mixture immediately after mixing.

The package leaflet of Ingelvac CircoFLEX should also be consulted before the administration of the mixed product.

WITHHOLDING PERIOD

Zero (0) days

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI):

Zero (0) days. Before using this product, confirm the current ESI from Boehringer Ingelheim Pty Limited on 1800 038 037 or the APVMA website (www.apvma.gov.au/residues)

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Additional user safety:

Take care to avoid self-injection.

Accidental self-administration may result in local bruising, pain and swelling. In the event of self-administration, seek medical attention and show the package leaflet or label to the Medical Practitioner.

Disposal

Dispose of empty container by wrapping in paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.

Storage

Store between 2-8°C. Do not freeze. Protect from light.

Not all pack sizes may be marketed.

APVMA Approval No.: 69810/61779

Boehringer Ingelheim Pty Limited

Animal Health Division
78 Waterloo Road
North Ryde NSW 2113
Australia

Ingelvac MycoFLEX® and Ingelvac CircoFLEX® are registered trademarks of Boehringer Ingelheim Vetmedica GmbH, used under license.



Boehringer Ingelheim

Revision 4
Operator hal
Date 29Jun15
Art # 1273285-00
SAP # 113688-001
Product Ingelvac
MycroFLEX
Type/Size Insert
Template # 671
Dimensions 5-5/8" x 8-3/4"
Fold 1-7/8" x 1-7/8"
PDF 29Jun15 hal

BLACK

142.875 mm x 222.25 mm

Art Number: 7 pt

Tradename: 14 pt

Text: 7.8 pt