

151603-01



FOR ANIMAL TREATMENT ONLY

Protech® C2i

Canine Coronavirus & Leptospira Copenhageni Killed Vaccine

COMPOSITION

Protech C2i Canine Coronavirus & Leptospira Copenhageni Killed Vaccine is prepared from virulent canine coronavirus (RP ≥ 1.62), blended with bacterins of *Leptospira interrogans* serovar Copenhageni (RP ≥ 1.0) together with a suitable adjuvant system.

Indication

For the immunisation of healthy dogs, 6 weeks of age or older to protect against disease caused by *Leptospira interrogans* serovar Copenhageni and aid in protection against disease caused by canine coronavirus.

This vaccine can be used for the prevention of clinical disease, however it may fail to prevent infection or organism shedding.

DIRECTIONS FOR USE

- Vaccinate only healthy dogs with normal temperatures.
- Dogs under treatment with immunosuppressive drugs should not be vaccinated.
- Studies to support the use of this product in pregnant bitches have not been conducted.
- Do not use chemicals or disinfectants to sterilise the skin, needles or syringes.
- A new sterile syringe and needle should always be used to administer this vaccine.

Precautions and side effects

Some animals may show transient post-vaccination pain, malaise and depression.

Allergic reactions such as anaphylaxis occur rarely and may require parenteral treatment with an antihistamine, a corticosteroid and/or adrenalin as appropriate.

Rarely, transient local site reactions after vaccination including swelling, pain and subcutaneous thickening may occur.

Dose: Withdraw the contents of a single dose vial (1 mL) aseptically into a sterile syringe and inject subcutaneously.

Protech C2i vaccine may be used as a diluent for Duramune Adult® C3 or C4, or Protech C3 or C4 vaccines.

Recommended primary vaccination program

In dogs previously unvaccinated with Protech C2i vaccine, two doses 2 to 4 weeks apart will protect against *Leptospira interrogans* serovar Copenhageni and canine coronavirus.

Revaccination

For protection against canine coronavirus and *Leptospira interrogans* serovar Copenhageni, annual revaccination with Protech C2i vaccine is recommended.

This vaccine has been assessed as providing at least 12 months protection. Many factors influence the effectiveness of vaccination and the need for revaccination. The vaccination program for an individual animal should be determined within a veterinary-client-patient relationship, taking all these factors into account.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

User Safety Information

Take care to avoid self-injection. Accidental self-administration may result in local bruising, pain and swelling. In the event of self-administration, seek medical attention and show the package leaflet or the label to the medical practitioner.

Disposal

Discarded syringes and needles should immediately be placed in a designated and appropriately labelled "sharps" container. Discarded vials should immediately be placed in a designated and appropriately labelled "biologicals" container.

Storage

Store between 2-8°C. Do not freeze. Protect from light.

Packaging

Supplied as 25 or 10 vials of killed vaccine. Not all pack sizes may be marketed.

Protech® and Duramune Adult® are registered trademarks of Boehringer Ingelheim Vetmedica, Inc. - used under license.

APVMA Approval No. 51488/60396

Boehringer Ingelheim Pty Limited
Animal Health Division
78 Waterloo Road
NORTH RYDE NSW 2113



Please recycle leaflet, label and tray only.

PC2iINST2

151603-01

Boehringer Ingelheim

Revision 4
Operator Ish
Date 09Jan14
Art # 151603-01
SAP #
Item # 14979W
Product Protech C2i
Type/Size Insert
Template # 534
Dimensions 3" x 7"
(76.2 x 177.8 mm)
Folds To 3" x 1-3/4"
PDF 10Jan14 Ish
Rev Level 01

BLACK

This file is 100% when printed on A4
(or A3, as applicable)

DIMENSIONS: 76.2 mm x 177.8 mm

FONT SIZES:

Tradename: 19 pt

Main Panel: 8 pt

ID#: 7 pt