

PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Ventipulmin[®]



Bronchodilator for Horses

ACTIVE CONSTITUENT:
CLENBUTEROL HYDROCHLORIDE 0.03 mg/mL

Properties

Ventipulmin[®] is a sympathomimetic amine with a high degree of selectivity for the beta-2 receptor sites in the body. These sites are primarily found in the lungs and uterus. This selectivity allows intense bronchodilating properties with minimum effect on the cardiovascular system.

Ventipulmin[®] also has a stabilising effect on mast cells, preventing release of histamine, serotonin, bradykinin and SRSA.

Ventipulmin[®] has a beneficial effect on mucociliary clearance by increasing the amount of mucus cleared by the mucociliary escalator.

Ventipulmin[®] dose rates for parenteral and oral forms of treatment are identical.

Indications

Bronchodilator for horses. For the treatment of all conditions where constriction of the airways occurs. Treatment of bronchospasm and airways obstruction associated with:
Bacterial infection – during and after the infective phase
Viral infection – during and after the infective phase
Bronchitis
Bronchopneumonia

Respiratory allergies – eg. to pollens, fungi, moulds and stable dust
Exercise induced bronchospasm
All conditions where narrowing of the small airways is a contributing factor to the problem.

DIRECTIONS FOR USE

Restraints

NOTTO BE USED in food producing species of animals.

Contraindications and Precautions
Ventipulmin[®] may inhibit the effect of prostaglandin F_{2a} and oxotocin. Use during pregnancy must be discontinued at term since uterine contractions may be abolished under its influence. This product is contraindicated for use with other beta adrenergic agents.

Dosage

Discard 28 days after first broaching the bottle.

1.25 mL per 50 kg bodyweight twice the first day by slow intravenous injection (or given intramuscular) followed by treatment with Ventipulmin[®] Oral Bronchodilator for horses for a minimum period of 5 days.

**MEAT WITHHOLDING PERIOD
(HORSES):
NOT TO BE USED in horses intended
for human consumption.**

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 131126; New Zealand 0800 764 766 (0800 POISON).

Disposal

Dispose of empty container by wrapping with paper and putting in garbage.

Presentation

50 mL injectable solution.

Storage

Store below 25°C (air conditioning)
Protect from direct sunlight.
Discard 28 days after first broaching the bottle.

Distributed by
Australia

**Boehringer Ingelheim Pty Limited
Animal Health Division
78 Waterloo Road
NORTH RYDE NSW 2113**

APVMA No. 35946/0905

New Zealand Information Withholding Period

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards. Horses producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 28 days of cessation of last treatment.

New Zealand

Boehringer Ingelheim (NZ) Ltd
Animal Health Division
Level 1, Unit 9
42 Ormiston Road, East Tamaki
Auckland

Restricted Veterinary Medicine A3891

See www.foodsafety.govt.nz
for registration conditions.